

# The Cosmic Dispersion Measure of Fast Radio Bursts using the EAGLE Simulations

Adam Batten

Swinburne University of Technology, Melbourne Australia

abatten@swin.edu.au



@adamjbatten

ANITA Workshop

2021-02-12

**ASTRO 3D**

SWIN  
BUR  
NE


Centre for  
Astrophysics and  
Supercomputing



# What is the Intergalactic Medium?



# What is the Intergalactic Medium?



Tumlinson et al. (2017)


# Why Do We Care About the Intergalactic Medium?

# Why Do We Care About the Intergalactic Medium?




1. The IGM contains most of the baryonic matter

# Why Do We Care About the Intergalactic Medium?



1. The IGM contains most of the baryonic matter



# Why Do We Care About the Intergalactic Medium?



1. The IGM contains most of the baryonic matter

2. Galaxies and the IGM evolve together

# Why Do We Care About the Intergalactic Medium?



1. The IGM contains most of the baryonic matter

2. Galaxies and the IGM evolve together

# Problems Observing the Intergalactic Medium

- **Density**  $\sim 1$  particle per cubic meter
- **Temperature**  $\sim 1 \times 10^6$  K


# Problems Observing the Intergalactic Medium

- Density  $\sim 1$  particle per cubic meter
  - Temperature  $\sim 1 \times 10^6$  K
- Lack of favourable UV/Optical transition lines. Hard to observe!!!

# Problems Observing the Intergalactic Medium

- Density  $\sim 1$  particle per cubic meter
- Temperature  $\sim 1 \times 10^6$  K
- ▶ Lack of favourable UV/Optical transition lines. Hard to observe!!!


**The Missing Baryon Problem:**  
 **$\sim 30\%$  of baryons at low redshift appear to be missing!**



Shull et al. (2012)


# How do Fast Radio Bursts (FRBs) help?

Lorimer et al. (2007)  
Petroff et al. (2019)



# How do Fast Radio Bursts (FRBs) help?

Lorimer et al. (2007)  
Petroff et al. (2019)




$$t = k\nu^{-2} \int_0^d n_e \, dl$$



# How do Fast Radio Bursts (FRBs) help?

Lorimer et al. (2007)  
Petroff et al. (2019)




$$t = k\nu^{-2} \int_0^d n_e \, dl$$

Electron  
number  
density



# How do Fast Radio Bursts (FRBs) help?

Lorimer et al. (2007)  
Petroff et al. (2019)




$$t = k \nu^{-2} \int_0^d n_e \, dl$$

Frequency


Electron number density

Diagram illustrating the dispersion measure (DM) equation. The equation is  $t = k \nu^{-2} \int_0^d n_e \, dl$ . A green arrow points upwards from the  $\nu^{-2}$  term to the word "Frequency". A pink arrow points downwards from the  $n_e$  term to the text "Electron number density".



# How do Fast Radio Bursts (FRBs) help?

Lorimer et al. (2007)  
Petroff et al. (2019)



Constant

$$t = k \nu^{-2}$$

Frequency


$$\int_0^d n_e \, dl$$

Electron  
number  
density



# How do Fast Radio Bursts (FRBs) help?

Lorimer et al. (2007)  
Petroff et al. (2019)




Constant

$$t = k \nu^{-2} \int_0^d n_e \, dl$$

Frequency

$$\text{DM} = \int_0^d n_e \, dl$$



$$DM = DM_{MW} + DM_{cosmic}(z) + \frac{DM_{Host}}{1+z}$$

$$\text{DM} = \text{DM}_{\text{MW}} + \text{DM}_{\text{cosmic}}(z) + \frac{\text{DM}_{\text{Host}}}{1+z}$$

Milky Way

↓

Host Galaxy/Local Environment

↓


$$\text{DM}_{\text{cosmic}}(z) = \text{DM}_{\text{IGM}}(z) + \text{DM}_{\text{CGM, Interlopers}}$$

↑


Intergalactic Medium

↑

The CGM of Galaxies along the line-of-sight



Ioka (2003) [Analytic]




Ioka (2003) [Analytic]


Inoue (2004) [Analytic]

Zhang (2018) [Analytic]

Macquart+(2020) [Analytic]





Ioka (2003)	[Analytic]
Inoue (2004)	[Analytic]
Zhang (2018)	[Analytic]
McQuinn (2014)	[Analytic+Hydro]
Dolag+(2015) [Hydro; Magneticum]	
Jaroszynski (2019) [Hydro; Illustris]	
Macquart+(2020)	[Analytic]



Ioka (2003)	[Analytic]
Inoue (2004)	[Analytic]
Zhang (2018)	[Analytic]
McQuinn (2014)	[Analytic+Hydro]
Dolag+(2015)	[Hydro; Magneticum]
Jaroszynski (2019)	[Hydro; Illustris]
Pol+(2019)	["Semi-Analytic"; MICE]
Macquart+(2020)	[Analytic]

# FRBs and the Missing Baryon Problem



# Batten et al. accepted (arxiv:2011.14547)

## The Cosmic Dispersion Measure in the EAGLE Simulations

Adam J. Batten,<sup>1,2</sup> Alan R. Duffy,<sup>1,2</sup> Natasha A. Wijers,<sup>3</sup> Vivek Gupta,<sup>1</sup> Chris Flynn,<sup>1</sup> Joop Schaye,<sup>3</sup> Emma Ryan-Weber<sup>1,2</sup>

<sup>1</sup>*Centre for Astrophysics and Supercomputing, Swinburne University of Technology, Melbourne, Victoria 3122, Australia*

<sup>2</sup>*ARC Centre of Excellence for All Sky Astrophysics in 3 Dimensions (ASTRO 3D)*


<sup>3</sup>*Sterrewacht, Leiden University, Niels Bohrweg 2, 2333 CA Leiden, The Netherlands*

Accepted XXX. Received YYY; in original form ZZZ

### ABSTRACT


The dispersion measure (DM) of fast radio bursts (FRBs) provides a unique way to probe ionised baryons in the intergalactic medium (IGM). Cosmological models with different parameters lead to different DM-redshift (DM –  $z$ ) relations. Additionally, the over/under-dense regions in the IGM and the circumgalactic medium of intervening galaxies lead to scatter around the mean DM –  $z$  relations. We have used the Evolution and Assembly of GaLaxies and their Environments (EAGLE) simulations to measure the mean DM –  $z$  relation and the scatter around it using over one billion lines-of-sight between redshifts  $0 < z < 3$ . We investigated two techniques to estimate line-of-sight DM: ‘pixel scrambling’ and ‘box transformations’. We find that using box transformations (a technique from the literature) causes strong correlations due to repeated replication of structure. Comparing a linear and non-linear model, we find that the non-linear model with cosmological parameters, provides a better fit to the DM –  $z$  relation. The differences between these models are the most significant at low redshifts ( $z < 0.5$ ). The scatter around the DM –  $z$  relation is highly asymmetric, especially at low redshift ( $z < 0.5$ ), and becomes more Gaussian as redshift approaches  $z \sim 3$ , the limit of this study. The increase in Gaussianity with redshift is indicative of the large scale structures that is better probed with longer lines-of-sight. The minimum simulation size suitable for investigations into the scatter around the DM –  $z$  relation is 100 comoving Mpc. The DM –  $z$  relation measured in EAGLE is available with an easy-to-use python interface in the open-source FRB redshift estimation package FRUITBAT.

**Key words:** intergalactic medium – hydrodynamics – methods: numerical – radio continuum: general



# EAGLE Simulations


Schaye et al. (2015), Crain et al (2015)



# EAGLE Simulations

Schaye et al. (2015), Crain et al (2015)


- Hydrodynamics + Nbody
- Large cosmological volume (100 cMpc)
- Redshift range ( $z \sim 127$  to  $z = 0$ )



# EAGLE Simulations

Schaye et al. (2015), Crain et al (2015)


- Hydrodynamics + Nbody
- Large cosmological volume (100 cMpc)
- Redshift range ( $z \sim 127$  to  $z = 0$ )
- Abundances for 11 different elements.
- HM12 UV Ionising Background
- Galactic Winds: Star formation & AGN



# EAGLE Simulations


Schaye et al. (2015), Crain et al (2015)

- Hydrodynamics + Nbody
- Large cosmological volume (100 cMpc)
- Redshift range ( $z \sim 127$  to  $z = 0$ )
- Abundances for 11 different elements.
- HM12 UV Ionising Background
- Galactic Winds: Star formation & AGN
- Resolution:  $\sim 0.7$  ckpc
- Particle Masses:  $\sim 10^6 M_\odot$



EAGLE Simulations

32,000




# EAGLE Simulations

- Divide cube into columns

32,000


32,000




# EAGLE Simulations

- Divide cube into columns
- Calculate column densities
  - Rahmati et al. (2013) (SS)
  - Wijers et al. (2019)

32,000




32,000




# EAGLE Simulations


- Divide cube into columns
- Calculate column densities
  - Rahmati et al. (2013) (SS)
  - Wijers et al. (2019)
- Convert column densities to units of  $\text{pc cm}^{-3}$






Local  
Observer




$z = 0$   
Observer










Increasing Redshift




Increasing Redshift












The  $\text{DM}_{\text{cosmic}}$  of most FRBs at low redshift appear to be  $2 - 3\sigma$  sigma above the mean.







# IGM Density Reconstruction Towards FRB 190608





Simha et. al 2020




FRB 190608

# IGM Density Reconstruction Towards FRB 190608




Simha et. al 2020

FRB 190608 intersects overdense IGM filaments along  
the line of sight!





FRB 190608

The  $\text{DM}_{\text{cosmic}}$  of most FRBs at low redshift appear to be  $2 - 3\sigma$  sigma above the mean.




# Mapping Foreground Large Scale Structure in FRB Fields



Credit: KG Lee

Using 2df on the AAT to follow up these FRBs and perform IGM reconstructions.





# Summary:



- The IGM is critical to understanding galaxy evolution and the Universe, but is difficult to observe at low redshift.
- Fast Radio Bursts provide a unique new way to probe the electron/baryon distribution in the IGM.
- **Paper Accepted: *The Cosmic Dispersion Measure in EAGLE***
  - arxiv:2011.14547
  - I used the EAGLE simulations to calculate DM-z relation and the scatter around it.
  - Large scatter around relation, with extremely skewed PDFs at low redshifts.
  - Most low redshift FRBs lie in the  $2-3\sigma$  confidence intervals.
  - Indicates intersection with IGM filaments, or possibly high host/source contributions.

# FRBs as Probes of Galaxy Feedback





# RefL0025N0752



# RefL0100N1504

